



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Since in forming the pair of elements one and only one element of a pair can be taken from the same set of planes, it follows that  $n(n-1)$  pairs can be taken from a same two sets of planes in which a corresponding pair of planes in the set are kept fixed and, therefore, in every possible way there can be formed

$$n(n-1)n_2 \text{ or } \frac{n^2 (n-1)^2}{2!}$$

by choosing in every possible manner the pair of planes to be kept fixed.

It should be noticed that the expansion passes directly from those terms involving only  $n-2$  elements of the principal diagonal plane to those involving them all. There would be no difficulty in stating at once the general law for the expansion of a cubical determinant in terms of the elements of the principal diagonal of the principal diagonal plane of the cubical determinant and their co-axial cubical minors.

---

## DEPARTMENTS.

---

### SOLUTIONS OF PROBLEMS.

---

#### ALGEBRA.

---

256. Proposed by THEODORE L. DeLAND, Treasury Department, Washington, D. C.

Three men, A, B, and C, rented a pasture for a fixed amount, each to pay per month in proportion to the stock pastured. During the first month A put in 3 horses and B and C each some horses, and B paid for the month \$6, but A and C each defaulted payment. During the next month each put in one more horse, and C paid for the month \$7.20, but A and B each defaulted payment. During the next month each put in one more horse, and A paid his bill for the month, \$5, but B and C each defaulted.

Required: (1) the rent of the pasture per month; (2) the number of horses B and C each put in during the first month; and (3) the amount A, B, and C, each, owed for the unpaid service.

I. Solution by S. A. COREY, Hiteman, Iowa.

Let  $a$ ,  $b$ ,  $c$ =rate per horse per month for the first, second, and third month, respectively. Let  $x$ ,  $y$ =number of horses put in the first month by B, and C, respectively. Let  $n=3+x+y$ =total number of horses in pasture first month. Let  $m$ =fixed monthly rental of pasture.

Then as A paid \$5 for the third month's rental when he had 5 horses in the pasture,  $c$ =\$1. As fewer horses were in the pasture the two preceding months the rate per horse per month was more than \$1 for the first and second months,

or both  $a$  and  $b > \$1$ . Then as  $x$  is an integer, as  $a > \$1$ , and as B paid but \$6 for the first month's service,  $x \leq 5$ . Similarly, as  $y$  is an integer,  $b > \$1$ , and as C paid but \$7.20 for the second month's service  $(y+1) \leq 7$ , or  $y \leq 6$ . Hence as  $n = 3 + x + y$ ,  $n \leq 14$ . But as  $c = m/(n+6) = 1$ ,  $m = (n+6)$ , whence  $m \leq \$20$ . But as not less than 5 horses were placed in the pasture the first month,  $m = \geq 11$ .

B had  $x$  horses in the pasture the first month, the rate then being  $m/n$ , or  $m/(m-6)$ , and paid \$6 for the service. Hence,  $xm/(m-6) = 36$ , or reducing and transposing,  $(6-x)m = 36$ , but as  $(6-x) = \text{integer}$ , and  $11 \leq m \leq 20$  (remembering that C had some horses in the pasture the first month, and that  $n = m - 6$ ).  $6 - x = 2$ ,  $m = \$18$ , whence  $x = 4$  and  $n = 12$ . But  $y = n - 3 - x = 5$ . As  $a = m/n = \$1.50$ ,  $b = m/(n+3) = \$1.20$ , and  $c = \$1$ , we readily find that A owed \$4.50 for the first, and \$4.80 for the second month's service = \$9.30 total. B owed \$6 for the second, and \$6 for the third month's service = \$12 total. C owed \$7.50 for the first, and \$7 for the third month's service = \$14.50 total.

II. Solution by REV. J. H. MEYER, S. J., Professor of Mathematics, College of the Sacred Heart, Augusta, Ga.; by M. R. BECK, Cleveland, Ohio, and by J. EDWARD SANDERS, Reinersville, Ohio.

Using  $r$  as the number of dollars in one month's rent;  $x$  and  $y$  as the number of horses B and C put in at first, respectively, we obtain

$$\frac{rx}{3+x+y} = 6, \text{ or } r = \frac{18+6x+6y}{x}. \quad (1)$$

$$\frac{r(y+1)}{6+x+y} = 7.20, \text{ or } r = \frac{43.2+7.2x+7.2y}{y+1}. \quad (2)$$

$$\frac{5r}{9+x+y} = 5, \text{ or } r = \frac{45+5x+5y}{5}. \quad (3)$$

Equating (1) and (3),  $90 - 15x + 30y = 5x^2 + 5xy$ ..... (4). Equating (2) and (3),  $171 + 31x - 14y = 5xy + 5y^2$ ..... (5). Adding (4) and (5),  $-5(x+y)^2 + 16(x+y) = -261$ ..... (6). Solving (6) for  $x+y$ ,  $x+y = 9$ ..... (7). Substituting (7) in (3), (2), (1),  $r = 18$ ,  $y = 5$ ,  $x = 4$ . For first month, \$1.50, for the second, \$1.20, and for the third, \$1.00 must be paid for each horse.

Hence, A owed \$9.30; B, \$12.00; C, \$14.50.

Also solved by O. L. Callecot, A. H. Holmes, L. E. Newcomb, J. Scheffer, G. B. M. Zerr, and the Proposer.

257. Proposed by L. E. NEWCOMB, Los Gatos, Cal.

Solve (1)  $x+y=10$ , (2)  $3x=\log_{10} y$ .

I. Solution by HENRY HEATON, Belfield, N. D.

We have  $3x = \log_{10}(10-x)$ , or  $10-x = 10^{3x}$ . If we suppose  $x = \frac{1}{3}$ , we obtain  $9\frac{2}{3} = 10$ . An error of  $\frac{1}{3}$ . If we suppose  $x = \frac{1}{4}$ , we get  $9.25 = 4.38$ . An error of 4.87. By position we obtain  $x = .328$ . Substituting this we get  $9.672 = 9.639$ . An error of  $-.043$ . Substituting .329 for  $x$  we get  $9.671 = 9.704$ . An error of